



URL: http://www.uni-jena.de/en/News/PM100708_exoplanet.pdf

The hunt for new Earths goes forward with a brand new planet-hunting technique

Astronomers found WASP-3c

A team of astronomers from Germany, Bulgaria and Poland have used a completely new technique to find an exotic extrasolar planet. The same approach is sensitive enough to find planets as small as the Earth in orbit around other stars. The group, led by Dr Gracjan Maciejewski of Jena University in Germany, used Transit Timing Variation to detect a planet with 15 times the mass of the Earth in the system WASP-3, 700 light years from the Sun in the constellation of Lyra. They publish their work in the journal *Monthly Notices of the Royal Astronomical Society*.

Transit Timing Variation (TTV) was suggested as a new technique for discovering planets a few years ago. Transits take place where a planet moves in front of the star it orbits, temporarily blocking some of the light from the star. So far this method has been used to detect a number of planets and is being deployed by the Kepler and Corot space missions in its search for planets similar to the Earth.

If a (typically large) planet is found, then the gravity of additional smaller planets will tug on the larger object, causing deviations in the regular cycle of transits. By comparing the deviations with extensive computer-based calculations, astronomers are able to deduce the makeup of the planetary system. For this search, the team used the 90-cm telescopes of the University Observatory Jena and the 60-cm telescope of the Rohzen National Astronomical Observatory in Bulgaria to study transits of WASP-3b, a large planet with 630 times the mass of the Earth.

"We detected periodic variations in the transit timing of WASP-3b. These variations can be explained by an additional planet in the system, with a mass of 15 Earth-mass (i.e. one Uranus mass) and a period of 3.75 days", said Dr Maciejewski. *"In line with international rules, we called this new planet WASP-3c".* This newly discovered planet is among the least massive planets known to date and also the least massive planet known orbiting a star which is more massive than our Sun. The host star WASP-3 is also younger than our Sun.

First detection by TTV

This is the first time that a new extra-solar planet has been discovered using this method. The new TTV approach is an indirect detection technique, such as the previously successful radial velocity and transit methods.

The discovery of the second, 15 Earth-mass planet makes the WASP-3 system very intriguing. The new planet appears to be trapped in an external orbit, twice as long as the orbit of the more massive planet. Such a configuration is probably a result of the early evolution of the system.

The TTV method is very attractive, because it is particularly sensitive to small perturbing planets, even down to the mass of the Earth. For example, an Earth-mass planet will pull on a typical gas giant planet orbiting close to its star and cause deviations in the timing of the larger objects' transits of typically a few seconds.

This is a big enough effect to be detected with relatively small 1-m diameter telescopes and discoveries can be followed up with larger instruments. The team are now using the 10-m Hobby-Eberly Telescope in Texas to study WASP-3c in more detail.

Contact:

Dr Gracjan Maciejewski / Prof. Dr Ralph Neuhäuser
Astrophysical Institute of Friedrich Schiller University Jena
D-07740 Jena
Phone: +49-3641-947524 / 947500
Mobile: +48-691-239648
Email: /

Further information:

The research appeared in "Transit timing variation in exoplanet WASP-3b", Maciejewski G., Dimitrov D., Neuhauser R., Niedzielski A., Raetz St., Ginski Ch., Adam Ch., Marka C., Moulla M., Mugrauer M., Monthly Notices of the Royal Astronomical Society, in press. A preprint of the paper can be found at <http://xxx.lanl.gov/abs/1006.1348>.

Meldung vom: 09.07.2010 10:00 Uhr